

GOES-19 SEISS EHIS Level 1b (L1b) Data Release
Provisional Data Quality
February 7, 2025
Read-Me for Data Users

The Peer Stakeholder - Product Validation Review (PS-PVR) for the GOES-19 Space Environment In-Situ Suite (SEISS) Energetic Heavy Ion Sensor (EHIS) L1b Provisional Maturity was held on February 7, 2025. As a result of this review, NOAA has confirmed that the GOES-19 EHIS L1b data are at Provisional Validation Maturity as of February 7, 2025. This Provisional declaration means that the data are ready for operational use, but the GOES-19 spacecraft is not yet deemed operational. Therefore, these data are to be accompanied by the "GOES-19 Preliminary, Non-Operational Data" caveat.

The EHIS L1b data product consists of 5-minute-cadence differential directional fluxes and associated systematic (instrumental) and statistical errors. Fluxes are produced for hydrogen (H) and helium (He); for the carbon-nitrogen-oxygen (CNO), neon-sulfur (Ne-S), and chlorine-nickel (Cl-Ni) mass groups; and for individual elements between beryllium and copper (Be-Cu) (but see below for a restriction on the beryllium and boron fluxes). EHIS has a single 60-degree (full cone angle) field-of-view directed radially outward from the Earth (toward zenith). The energy range is nominally 10-200 MeV/nucleon for hydrogen (protons) and helium (alpha particles), divided into five energy channels. The energy range increases with atomic number (Z) since the stopping power in silicon is the same for all species in each energy channel. Outside of solar energetic particle (SEP) events, EHIS observes backgrounds, some of which are consistent with galactic cosmic ray (GCR) flux levels.

The H and He fluxes are derived directly from coincidence rates (3-second cadence in the raw Level 0 data), as with SGPS, and can be averaged over longer periods to improve the counting statistics. However, the heavy ion fluxes are derived using a maximum likelihood (ML) fit to a histogram of Z values determined on-orbit (sum of five 1-minute cadence histograms) using the angle-detecting inclined sensor (ADIS) system incorporated into the EHIS telescope. While this ML fit is necessary for meeting requirements in the presence of very sparse heavy ion count rates, it limits the utility of the L1b data in post-processing.

From the Ground Processing Algorithm Document for the GOES-R Space Environment In-Situ Suite (SEISS), Rev. F (p. 76): "EHIS data is accumulated over 3 second and 1 minute intervals. Data products for longer periods of time must be added together from the raw data and processed as shown below. Taking fluxes from five [or] 1-minute periods (particularly upper limits) and simply averaging them to obtain fluxes for a longer period, is not valid and EHIS reporting requirements will not be met." Moreover, when, in the L1b data, the lower one-sigma statistical error is equal to the mean value, only an upper limit exists (mean plus upper one-sigma statistical error). (In the L1b files, the mean fluxes are contained in the variable 'BeCu5MinuteDifferentialFluxes', and the lower and upper statistical errors are contained in the variable 'BeCu5MinuteDifferentialFluxStatErrorsBounds'.) As a result, derivation of heavy ion fluxes for periods longer than 5 minutes requires reprocessing from Level 0 raw data. This is a limitation that is independent of the maturity of the product.

Provisional validation means:

- Validation activities are ongoing and the general research community is now encouraged to participate.

- Severe algorithm anomalies are identified and under analysis. Solutions to anomalies are in development and testing.
- Incremental product improvements may still be occurring.
- Product performance has been demonstrated through analysis of a small number of independent measurements by another instrument on the same observatory, or by other observatories.
- Product analysis is sufficient to establish product performance relative to expectations (Performance Baseline).
- Documentation of product performance exists that includes recommended remediation strategies for all anomalies and weaknesses. Any algorithm changes associated with severe anomalies have been documented, implemented, and tested.
- Testing has been fully documented.
- Product is ready for operational use and for use in comprehensive calibration/validation activities and product optimization.

Users of the GOES-19 EHIS L1b data bear responsibility for inspecting the data and understanding the known caveats prior to use. Below is the list of caveats that have been identified and are under analysis. Solutions are in development and testing.

1. Current understanding of GOES-19 EHIS performance is based on analysis of the SEP event of 09-11 October 2024, during which EHIS observed some significant heavy ion fluxes (carbon, nitrogen, and oxygen in the lowest energy channel). Up to the date of the provisional PS-PVR, this was the largest SEP event of Solar Cycle 25 in terms of >10 MeV protons, though not in terms of the heavy ion energies observed by EHIS.
2. EHIS hydrogen and helium fluxes are low relative to SGPS. The root cause of this discrepancy has not been determined. In general, NOAA recommends that the SGPS proton (hydrogen) and alpha particle (helium) fluxes be used instead of the EHIS helium channels. SGPS produces hydrogen and helium fluxes over a wider energy range and in more energy channels than EHIS. However, since EHIS has an order-of-magnitude greater event-to-background ratio than SGPS at the lowest energies, studies of weak events may benefit from using EHIS data, keeping in mind the absolute calibration differences.
3. Since May 2023, the fluxes in the two lowest-energy helium channels in the L1b EHIS files from all GOES-R series satellites are replaced with interpolated SGPS data. The decision to do this was based primarily on the fact that EHIS helium fluxes are low relative to SGPS, without considering the unique strengths of EHIS data, in particular its greater event-to-background ratio and its zenith look direction. This look direction is orthogonal to the eastward and westward look directions of the two SGPS units on each satellite and therefore provides additional scientific information on anisotropies in the solar particle fluxes. NOAA recommends against use of the two lowest-energy EHIS helium channels from the L1b files - instead, please use the SGPS alpha particle fluxes in the Level 2 five-minute-averages files, which have better energy resolution.
4. As of February 7, 2025, there are two residual errors related to this replacement of EHIS helium fluxes with interpolated SGPS fluxes. First, the energies of the two lowest-energy helium channels are recorded as zero in the files. Second, when a satellite is flying inverted (as GOES-19 will for approximately six

months starting in late March 2025), the fluxes in the two lowest-energy helium channels are recorded as zero. These errors are expected to be fixed during Spring 2025.

5. No EHIS L1b data processed prior to declaration of Provisional Maturity (e.g., those available from CLASS) should be used, owing to an important table update prior to Provisional Maturity.
6. Absolute calibrations are still being refined. For example, a temperature sensitivity that may impart a significant diurnal variation onto the observed proton (hydrogen) fluxes remains to be quantified.
7. Lithium, beryllium and boron ions of solar origin are never observed, being destroyed in the solar interior. Therefore, L1b Be and B fluxes are replaced with fill values because they should never be used. (Lithium fluxes are not within the reported measurement range.) It is believed that counts at these histogram locations are due to slow helium nuclei being counted as heavy ions.
8. Outside of SEP events, EHIS observes backgrounds, some of which may be from GCR fluxes. Under these conditions, the L1b fluxes are not accurate, since the processing uses geometrical factors and energy bandwidths derived for SEP spectra. Steep SEP spectra emphasize the lower-energy ends of the energy response functions while flatter GCR spectra more strongly emphasize the higher-energy tails of the response functions.
9. As described above, time-averaging L1b heavy ion fluxes, particularly those that are upper limits, does not result in improved accuracy and therefore should not be performed.

Contact for further information: OSPO User Services at SPSD.UserServices@noaa.gov

NCEI contacts for specific information on the EHIS L1b data:

Juan Rodriguez juan.rodriguez@noaa.gov

Brian Kress brian.kress@noaa.gov

Pamela Wyatt pamela.wyatt@noaa.gov

NCEI website for GOES-R Space Weather data (provides daily aggregations of EHIS L1b data):

<https://www.ngdc.noaa.gov/stp/satellite/goes-r.html>